

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Condor®

Selective
HERBICIDE

ACTIVE CONSTITUENTS:
375 g/L MCPA present as the potassium salt
10 g/L PYRAFLUFEN-ETHYL

GROUP	GI	HERBICIDE
-------	-----------	-----------

For the control of annual broadleaf weeds in wheat and oats as per the Directions For Use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.

Contents: 5L, 10L, 20L, 110L

APVMA Approval No: 86923 / 121811

SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcaml Pacific Australia Pty Ltd

CONDOR SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When opening the container and preparing spray, and applying by boom spray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

Batch Number:

Date of Manufacture:

Not a Dangerous Good for transport by road or rail according to the Australian Code for the Transport of Dangerous Goods by Road and Rail. Limit 500L

For specialist advice in emergency only, call 1800 033 111 all hours, Australia wide



G.H.S. STATEMENTS

Hazard H302+H312+H332: Harmful if swallowed, in contact with skin or if inhaled.

Prevention P261 Avoid breathing mist, vapours, spray. P270: Do not eat, drink or smoke when using this product. P271: Use only outdoors or in a well-ventilated area. P280: Wear protective gloves / protective clothing / eye protection / face protection.

Response P301 + P330 + P331: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. P302+P352: IF ON SKIN: Wash with plenty of soap and water. P304+P340: IF INHALED: Remove person to fresh air and keep at rest in a position comfortable for breathing. P312: Call a POISON CENTRE or doctor/physician if you feel unwell.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Condor®

Selective

HERBICIDE

ACTIVE CONSTITUENTS

375 g/L MCPA present as the potassium salt
10 g/L PYRAFLUFEN-ETHYL

GROUP	GI	HERBICIDE
-------	-----------	-----------

For the control of annual broadleaf weeds in wheat and oats as per the Directions For Use.

THIS LEAFLET IS PART OF THE LABEL

APVMA Approval No: 86923/121811

SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademarks of Sipcaml Pacific Australia Pty Ltd

DIRECTIONS FOR USE: All States

RESTRAINTS:

DO NOT apply by aircraft.

DO NOT apply where the slope exceeds 4%.

DO NOT tank mix Condor Selective Herbicide with any adjuvant, wetting agent, crop oil, insecticide, fungicide, trace element, fertiliser or other herbicide unless specified in the DIRECTIONS FOR USE and Crop Safety sections of the label.

DO NOT apply if rain is expected within 6 hours of application.

DO NOT apply to weeds suffering from stress caused by conditions such as frost, drought or soil water logging.

SPRAY DRIFT RESTRAINTS

DO NOT apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour, as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site. Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application, and must **KEEP** this record for at least 2 years. The spray application details that must be recorded are: **1** date with start and finish times of application **2** location address and paddock(s) sprayed **3** full name of this product **4** amount of product used per hectare and number of hectares applied to **5** crop or situation and weed or pest **6** wind speed and direction during application **7** air temperature and relative humidity during application **8** nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application **9** name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers within **100 metres** downwind from the application area.

DO NOT apply if there are sensitive crops, gardens, landscaping vegetation, protected native vegetation or protected animal habitat within **140 metres** downwind from the application area.

Crop	Weeds Controlled	Rate	Crop Stage	Weed stage	Critical Comments
Wheat, oats	Bedstraw (<i>Galium tricornutum</i>), Bifora (<i>Bifora testiculata</i>), Capeweed	0.8 L/Ha	Minimum 2 leaf	2-4 leaf	Apply as a post-emergence treatment to actively growing weeds. The addition of a non-ionic surfactant may improve weed control. Add 100mL/100L of a 1000 g/L alcohol alkoxyate non-ionic surfactant eg BS1000*. Under favourable growing conditions some weed regrowth may occur. A follow up application of a suitable herbicide may be required as part of a good weed management strategy.
	(<i>Arctotheca calendula</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Long storks bill (<i>Erodium botrys</i>), Marshmallow (<i>Malva parviflora</i>)	1.0 L/Ha	Minimum 3 leaf to end of tillering	4-6 leaf	
	Prickly lettuce (<i>Lactuca serriola</i>), Volunteer canola (<i>Brassica napus</i>), Volunteer lupin (<i>Lupinus</i> sp), Wild turnip (<i>Brassica tournefortii</i>)	1.6 L/Ha	Minimum 3 leaf to end of tillering	6-8 leaf	
	Wild radish (<i>Raphanus raphanistrum</i>)				

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING:

Wheat, Oats: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Condor Selective Herbicide is an early post-emergence contact herbicide with rapid foliar uptake. Herbicide effects on susceptible weeds can occur within 1 to 7 days of application. Subsequent germinations will not be controlled.

APPLICATION

Prior to applying Condor Selective Herbicide, clean the spray tank to remove any other chemical, wetters or adjuvants remaining from previous spray operations otherwise crop injury may result. Apply as a broadcast application. Use conventional boom sprayers with either mechanical or bypass agitation. Spray equipment should be properly calibrated to ensure correct application. Use a spray volume of 70 to 150 litres per hectare. This is particularly important on hard-to-control weeds. Use the higher volume if weed infestation is heavy or the crop cover is dense. The best application conditions are when soil is moist, weather fine and rain unlikely within 6 hours.

RESISTANT WEEDS WARNING

GROUP G I HERBICIDE

Condor Selective Herbicide is a member of the phenylpyrazole and phenoxy group of herbicides. Its mode of action is through a process of membrane disruption, which is initiated by the inhibition of the enzyme protoporphyrinogen oxidase and disruption of plant cell growth. This inhibition interferes with the chlorophyll biosynthetic pathway. For weed resistance management, Condor Selective Herbicide is a Group G and a Group I herbicide. Some naturally occurring weed biotypes resistant to Condor Selective Herbicide and other herbicides that inhibit the enzyme protoporphyrinogen oxidase or disrupt plant cell growth may exist through normal genetic variability in any weed population and increase if these herbicides are used repeatedly. These resistant weeds will not be controlled by Condor Selective Herbicide or other herbicides that inhibit the enzyme protoporphyrinogen oxidase or disrupt plant cell growth. Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Condor Selective Herbicide or other herbicides that inhibit the enzyme protoporphyrinogen oxidase or disrupt plant cell growth.

CROP SAFETY

DO NOT tank mix Condor Selective Herbicide with any adjuvant, wetting agent, crop oil, insecticide, fungicide, trace element, fertiliser or other herbicide unless specified in the DIRECTIONS FOR USE and Crop Safety sections of the label. **DO NOT** mix Condor Selective Herbicide mixtures with selective grass herbicides as grass weed control will be significantly reduced and excessive crop injury may occur. Instead, allow a 10 to 14 day interval between spraying Condor Selective Herbicide mixtures and grass herbicide applications.

Cereals

Some herbicidal symptoms may appear on wheat in the form of leaf spotting due to environmental conditions. However the crop recovers quickly usually with 2-3 weeks of treatment.

COMPATIBILITY:

Condor is compatible with Esteem® (600 g/Kg metsulfuron-methyl), Sipgran® (750 g/kg triasulfuron), Sylon® 750 SG (750 g/kg clopyralid), Volley® SG (700 g/Kg dicamba), Aspect (500g/L diflufenican) and certain trace elements. Condor may be mixed in the spray tank with one of the following insecticides or fungicides at label rates: Rover (400 g/L dimethoate), Alphasip Duo 100 (100 g/L alpha-cypermethrin), Le-mat (290 g/L omethoate) or Laguna (430 g/L tebuconazole). Do not apply metsulfuron methyl tank mixes on oats.

In all cases, do not add non-ionic surfactant when tank mixing with other products.

SPRAYER CLEAN-OUT

Do not allow the spray solution to dry in the application equipment. After application and before using the sprayer equipment for any other applications, the sprayer must be thoroughly cleaned. Immediately following application, clean all equipment thoroughly with detergent or a spray tank cleaner and water as described below. Should residues of Condor Selective Herbicide remain in inadequately cleaned equipment, they may be released in subsequent applications and cause injury to crops.

1. Drain sprayer tank, hoses and spray boom and thoroughly rinse with clean water the inside of the spray tank, sprayer hoses, boom, and nozzles to remove any sediment or residues.
2. Fill the tank with clean water, add an alkaline detergent or boom cleaner (follow manufacturer's directions for use). Fill tank to capacity and operate the sprayer with agitation for 15 minutes to flush hoses, boom and nozzles.
3. Drain the sprayer tank, lines, and booms. Rinse the tank with clean water and flush through the hoses, boom and nozzles. Remove and clean spray nozzles, tips and screens.

PRECAUTIONS

Re-entry

Do not allow entry into treated areas until spray has dried. If prior entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NONTARGET PLANTS

Do NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK

Certain plants such as Caltrop, Capeweed, Paterson's Curse and Variegated Thistle may be naturally toxic to stock when eaten in large quantities under certain conditions. Where plants are known to be toxic, grazing should be delayed until complete browning of treated plants has occurred. Do not graze horses or pigs on Paterson's Curse.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. Do NOT contaminate wetlands or watercourses with this product or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When opening the container and preparing spray, and applying by boom spray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.